

Best Management Practices for Colorado Corn

Primary Authors: Troy Bauder
Department of Soil and Crop Sciences
Colorado State University

Reagan Waskom
Department of Soil and Crop Sciences
Colorado State University

Contributing Authors: Joel Schneekloth
Colorado State University Cooperative Extension

Jerry Alldredge
Colorado State University Cooperative Extension

Technical Writing and Support: Marjorie Nockels Ortiz
Layout: Debbie Fields

Graphic Design: Nancy Reick, Kendall Printing

Funded by the Colorado Corn Growers Association/Colorado Corn Administrative Committee through a grant by the Colorado Department of Public Health and the Environment through a Section 319 Nonpoint Source Education Grant. Additional funding and support provided by the Agricultural Chemicals and Ground Water Protection Program at the Colorado Department of Agriculture.

Special acknowledgments to
the following reviewers:

Bruce Bosley, Colorado State University Cooperative Extension
Bill Brown, Colorado State University
Grant Cardon, Colorado State University
Wayne Cooley, Colorado State University Cooperative Extension
Bill Curran, Pioneer Hi-Bred International, Inc.
Ron Meyer, Colorado State University Cooperative Extension
Frank Peairs, Colorado State University
Calvin Pearson, Colorado State University
Gary Peterson, Colorado State University
Dwayne Westfall, Colorado State University
Phil Westra, Colorado State University

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Milan A. Rewerts, Director of Cooperative Extension, Colorado State University, Fort Collins, Colorado. Cooperative Extension programs are available to all without discrimination. To simplify terminology, trade names of products and equipment are occasionally used. No endorsement of products mentioned is intended nor is criticism implied of products not mentioned.

Published by Colorado State University Cooperative Extension in Cooperation with the Colorado Corn Growers Association/
Colorado Corn Administrative Committee.
Colorado State University Cooperative Extension Bulletin XCM574A. February 2003.

Nitrogen

Nitrogen (N) is the essential nutrient that most frequently limits corn yields in Colorado. Likewise, corn requires more added N from fertilizer or other sources than any other nutrient. Approximately 1.2 lbs N is required to produce one bushel of corn. Nitrogen management is also closely tied to irrigation management since N is mobile in the soil as nitrate (NO_3^-).

Good N management in corn involves making sure the crop has the right amount of available N at the right time. Therefore, timing of application is as important as amount, in certain situations.

Soil testing and crediting other N sources are the base practices for sound N management. Other tools that may increase N efficiency are tissue testing, irrigation water and manure analysis, and pre-sidedress soil nitrate testing (PSNT).

Nitrogen transformation in soil




Figure 16. The nitrogen cycle in soil. Nitrogen is not a stable element in the environment and can exist in many forms.

- NO_3^- and NH_4^+ become available from sources other than fertilizer and manure, such as crop residues and soil organic matter, legume fixation and atmospheric deposition.
- Added N fertilizer can become unavailable to corn through leaching, loss to gaseous forms (volatilization and denitrification), or conversion to organic N forms.



Nitrogen deficiency in corn. Appearing first on older leaves, yellowing starts at leaf tip and moves towards the base in a "V" pattern down the midrib.

Nitrogen in corn

- Plant takes up ammonium (NH_4^+) and nitrate (NO_3^-) forms.
- Nitrogen is mobile within plant & will move from older to younger tissues.
- NO_3^- is very water soluble and moves easily (leaches) in soil.
- NH_4^+ is adsorbed to soil particles and does not readily leach.

Soil Fertility

Good N management programs begin with a representative soil sample.

- Sample to a minimum of 2 feet deep for best results.
- Take 15 to 20 cores from a uniform area (no more than 40 acres).
- Base yield expectation on the average yield history for each field (5 years), plus 5%.

BMP

Identify fields with a high potential for leaching or runoff and employ all appropriate BMPs on these fields to reduce contamination of water supplies.

Determining correct nitrogen application rates

Each bushel of corn harvested contains 0.80 to 0.90 pounds of N and each ton of silage contains approximately 7.7 pounds of N per ton silage (35% DM). However, research as shown that the following equations closely predict corn N requirements in most situations. Since the gross N requirement is based on yield goal, use an accurate yield history when determining N rate.

Use the CSU N requirement calculation below to compare recommendations from soil testing laboratories. If the recommendations vary greatly, ask the lab to explain their procedures for calculating N requirements.

Nitrogen fertilizer requirement =

1. Total N required for yield expectation:

Grain: Total N needed = (yield expectation, bu x 1.2) + 35

Silage: Total N needed = (yield expectation (wet weight basis), tons x 7.5) + 35

2. Subtract soil organic matter from total N required:

Organic matter credit = (grain yield goal x 0.14) x % organic matter

Organic matter credit = (silage yield goal x 0.85) x % organic matter

3. Subtract residual soil NO₃-N, in the top 2 feet of soil, from total N required:

Soil NO₃-N credit = 8 x average soil NO₃-N

4. Subtract manure credit (see page 56, Table 8)

5. Subtract irrigation water credit (see Irrigation Nitrate Credit box, page 55)

= Irrigation water NO₃-N ppm x 0.223 lbs. N/acre inch x inches irrigation water

6. Subtract legume credit (if any, see Table 7)

Example fertilizer N calculation

200 bu yield goal

No manure or legume

1.5% OM

12 ppm NO₃-N in soil

15 ppm NO₃-N irrigation water

Total N needed = (200 bu x 1.2) + 35	= 275 lb N/A
organic matter credit: (200 bu x 0.14) x 1.5%	= -42
residual soil credit: 8 x 12 ppm NO ₃ -N	= -96
irrigation water credit: 1 ppm NO ₃ -N x 0.23 lb x 16" H ₂ O	= -55
Net N fertilizer required:	= 82 lbs N/A

Soil Fertility

Table 6. How to calculate average NO₃-N content across soil depths.

Soil layer sampled	Sample thickness	Measured NO ₃ -N	Calculations
-----inches-----		-----ppm-----	
0 - 8	8	20	8 x 20 = 160
8 - 24	16	8	16 x 8 = 128
			288/24" = 12ppm

How to calculate irrigation water nitrate credit

1. Determine the nitrate-nitrogen (ppm NO₃-N) content of the irrigation well water by field test kits or laboratory analysis. Initially, sample water twice a year to account for seasonal variability, and then annually.
2. Estimate the amount of water to credit. Most N uptake in corn occurs by milk stage (R3). Therefore, credit water applied before mid-August. Reasonable water credits in Colorado are: Northeast Colorado, 15 inches; Central High Plains, 17 inches; Southern High Plains and Arkansas Valley, 19 inches. Use the following equation to determine credit:

$$\text{NO}_3\text{-N ppm} \times 0.23 \times \text{H}_2\text{O inches} = \text{N credit (lbs/acre)}$$

* an acre-inch of water contains 0.23 lbs of N for each ppm of NO₃-N

* 1 ppm = 1 mg/L

Legume Credit

Table 7. Nitrogen credits for previous legume crop.

Legume Crop	Stand Quality	lb N/A credit*
Alfalfa	60-100% (4-6 plants per sq. ft.)	100 - 140
	30 - 60% (3-4 plants per sq. ft.)	70 - 100
	0 - 30% (0-3 plants per sq. ft.)	0 - 70
Dry beans		25

*Based upon a 3rd year stand or older.




Figure 17. Crediting nitrate is a BMP that protects ground water.

- Groundwater in irrigated areas along the South Platte and Arkansas Rivers frequently contains greater than 10 ppm NO₃-N. That amounts to over 27 lbs/ acre foot of water applied.
- Crediting irrigation water NO₃-N towards a N requirement can help remove nitrate from groundwater and lowers fertilizer costs.

Soil Fertility

Table 8. Approximate N credits for various solid manure sources in Colorado.

Manure	% H ₂ O	lb N/ton credit*	
		---dry---	---as is---
Beef	32	13	9
Dairy	46	9	5
Swine	82	22	4

* N credit for 1st crop year following application. For the second and third years use 1/2 and 1/4 of the first year N credits, respectively.

For more information, see CSU Bulletin 568A, BMP's for Manure Management.

BMP
 Incorporate manure as soon as possible after application to minimize volatilization losses, reduce odor, and prevent runoff. Phosphorus fertilizer or manure should not be applied to soils testing 'very high' for soil P. Limit solid manure application on frozen or saturated ground to fields not subject to runoff. Liquid effluent should not be applied to frozen or saturated ground. Plant permanent vegetation strips around the perimeter of surface water and erosive fields to catch and filter nutrients and sediments in surface runoff.

Manure

Livestock manure is an excellent nutrient source for corn production. Although manure is typically applied to meet nitrogen requirements, it also contains significant amounts of phosphorus, potassium, zinc and iron. Manure also improves soil tilth, water holding capacity, and infiltration.

While manure offers many benefits for soil health and fertility, it also poses risks. Continuous over-application of manure can result in increased salinity, degradation of surface water from phosphorus runoff, or groundwater contamination by nitrate. Growers should always apply manure based upon crop nutrient need determined by soil testing and reasonable yield goals. Use the Colorado P-index to determine appropriate application rates where soil test P is high or runoff may occur.

Table 9. Approximate manure rates needed for desired N application rate. Use the formula on page 54 to determine the N requirements and match to the desired N application in the left hand column.

Nitrogen content of manure						
Desired N application:	----- lbs of total N per ton manure (fresh weight basis) -----					
	8	13*	18	23**	28	28
(N lbs/acre)	Tons of manure per acre to apply for desired nitrogen application					
60	19	12	8	7	5	5
80	25	15	11	9	7	6
100	31	19	14	11	9	8
120	38	23	17	13	11	9
140	44	27	19	15	13	11
160	50	31	22	17	14	12
180	56	35	25	20	16	14
200	63	38	28	22	18	15

Tons to apply = $\frac{\text{Desired N application}}{(\text{total N content} \times 0.4)}$

*Average N content for Dairy manure in Colorado (wet weight)

** Average N content for beef manure in Colorado (wet weight)

Source: Cattle Manure Application Rates, CSU Extension Fact Sheet 0.561



Manure application